

META-ANALYSIS

Effects of intermittent overload doses of oral vitamin D₃ on serum 25(OH)D concentrations and the incidence rates of fractures, falls, and mortality in elderly individuals: A systematic review and meta-analysis

Xiaoyang Tao ¹, Wupeng Yang ¹, Qinxin Zhang ¹, Yongjiang Wang ¹, Feng Gao¹, Yuehan Wang ², Tingxin Zhang ¹, Hao Liu ^{1*}, and Jindong Chen ^{3*}

Vitamin D is commonly used to prevent and treat osteoporosis, with studies indicating its potential to reduce fractures, falls, and mortality. However, meta-analyses present inconsistent findings regarding its efficacy, particularly reflecting significant variability in data and outcomes related to various dosing regimens. In this meta-analysis, we assessed the impact of high-dose intermittent oral administration of vitamin D₃ on serum 25(OH)D levels, fractures, falls, and mortality among elderly individuals. We included 14 randomized controlled trials (RCTs) and employed Review Manager 5.4 for statistical analysis. Our findings indicate that intermittent monthly administration of vitamin D₃ (over 800 IU per day) significantly raised serum 25(OH)D levels at all timepoints after six months, maintaining levels above 75 nmol/L throughout the year. This regimen showed no increase in all-cause mortality, with a risk ratio (RR) (95% confidence interval [CI]) of 0.95 (0.87–1.04). Likewise, it did not significantly reduce the risks of falls and fractures, with RRs of 1.02 (0.98–1.05) and 0.95 (0.87–1.04), respectively. Although one-year intermittent administration significantly increased the concentration of 25(OH)D in serum, further research is needed to determine if this method would increase the incidence of falls. Therefore, it is not recommended at this stage due to the lack of demonstrated safety in additional relevant RCTs. This study had been registered at PROSPERO (CRD42022363229).

Keywords: Vitamin D₃, 25(OH)D, oral, load dose, intermittent, elderly.

Introduction

Vitamin D is not only widely used in the treatment of osteoporosis in middle-aged and elderly individuals but also has possible protective effects against cancer, infection, cardiovascular disease, and other diseases; therefore, it has wider indications [1–4]. It is generally considered that the current suitable concentration of 25(OH)D is between 30 and 11.94 ng/mL [5–7]. A lower dose (400 IU/day) has little effect on serum 25(OH)D concentrations, while 800 IU/day is the most commonly prescribed dosage [8]. In nursing home (NH) patients with severe 25(OH)D deficiency, an individually calculated cholecalciferol loading dose may be superior to a cholecalciferol 800 IU daily dose in rapidly normalizing vitamin D levels. This suggested that higher doses of vitamin D may be more rapid and effective in increasing serum 25(OH)D concentrations [9].

However, Dawson-Hughes and Harris also hypothesized that a 500,000 IU dose may trigger a “short-term

protective” reaction in which CYP24 (25-hydroxyvitamin D-24-hydroxylase), the enzyme that catalyzes 1,25(OH)₂D, is regulated, resulting in reduced serum and tissue levels of 1,25(OH)₂D [10]. This hypothesis was consistent with results from an animal study [11]. The randomized controlled trial (RCT) conducted by Glendening [12] showed that there was no statistically significant difference in the mean serum 25(OH)D levels between the experimental and control groups after nine months. Therefore, it is necessary to investigate the effect and safety of high-dose, intermittent oral vitamin D₃.

In a previous meta-analysis, two different drugs, vitamin D₂ and vitamin D₃, were combined [13–17]. However, the results were inconsistent or heterogeneous. From a pharmacodynamic perspective, vitamin D₃ has a greater ability to not only increase but also sustain higher serum 25(OH)D concentrations over time compared to vitamin D₂ [18]. In addition, considering the reduced autonomy of elderly individuals, the compliance of daily administration was worse compared with intermittent

¹Orthopedics Department, Ordos Central Hospital, Ordos, China; ²Ordos No.1 Middle School, Ordos, China; ³Ordos School of Clinical Medicine, Inner Mongolia Medical University, Ordos, China.

*Correspondence to Jindong Chen: chenjindong970416@163.com; Hao Liu: 31349015@qq.com

Associate editor: Gulali Aktas

DOI: 10.17305/bb.2024.10449

© 2024 Tao et al. This article is available under a Creative Commons License (Attribution 4.0 International, as described at <https://creativecommons.org/licenses/by/4.0/>).




Figure 1. Risk of bias assessment. Judgments for each risk of bias item presented as percentages across all included studies.

oral administration, resulting in more cost [19], so a high-dose and intermittent oral vitamin D₃ regimen may be more suitable for elderly individuals. Therefore, the purpose of this study was to investigate the clinical effects and safety profile of vitamin D₃ with a meta-analysis conducted under the specific conditions of loading dose and intermittent oral administration.

Materials and methods

Meta-analysis of randomized controlled trials (RCTs)

RCTs were eligible for inclusion if they met the following criteria: 1) RCTs comparing vitamin D₃ alone or in a combination with a placebo or a low dose (less than 400 IU per day); 2) The duration of the RCT was over 6 months, with the interval between doses more than 1 month, and each administration was a large dose (equivalent to more than 800 IU per day); 3) The average age of the participants was greater than 60 years old; 4) The mode of administration was limited to oral administration; and 5) The baseline serum 25(OH)D concentration of the included participants was greater than 30 nmol/L. The number of participants with one or more falls, fractures, and deaths were reported separately for the vitamin D treatment group and the control group.

Dosages were categorized as follows: 1) Low dose: Less than 400 IU per day; 2) Medium dose: Between 400 and 800 IU per day; and 3) High dose: Greater than 800 IU per day (and single dose greater than 40,000 IU).

Exclusion criteria were as follows: 1) RCTs with vitamin D₂ or bisphosphonates; 2) RCTs that used active vitamin D, which requires monitoring for hypercalcemia, with much higher costs, thereby limiting their public health applicability; 3) Studies including patients with diseases that may lead to a significant decline in autonomy or motor stability, such as Parkinson's disease, cerebral infarction, epilepsy, and other diseases; and 4) Studies that used intramuscular injections or intravenous administration.

Data extraction and quality assessment

This study was carried out independently by two researchers between October 2022 and January 2023, according to the Preferred Reporting Items for Systematic Reviews and

Meta-analysis (PRISMA) guidelines, and possible bias was assessed. PubMed, EMBASE, Cochrane Library, and other RCT databases were searched from database inception until January 30, 2023. We performed categorical analysis, heterogeneity checks, publication bias analysis, and subgroup analysis. The following data were extracted from the RCTs: year of publication, study design, sample size, duration of the intervention, percentage of women, number of total falls, fractures and deaths, serum 25(OH)D concentrations at different timepoints, and dosage and frequency of vitamin D administration. The authors of the included RCTs were contacted via e-mail for incomplete data. Some missing data were also derived from other previous analyses if the authors were unreachable. Quality assessment was performed by two independent researchers using the Cochrane Collaboration tool (Figure 1). We began our literature search in early 2024 while conducting the literature review. The risk of bias included in the literature was not high.

Data synthesis and statistical analysis

The main outcome was the serum 25(OH)D concentration at different periods, followed by the incidence of fractures, falls, and death. Because the 25(OH)D concentration results of the combined RCTs were in different units (nmol/L or ng/mL), the results for the continuous variable were calculated using the standardized mean difference method. The Mantel-Haenszel method was used to calculate risk ratios (RRs) and their 95% confidence intervals (CIs). The I^2 statistics or L'Abbe plots were used to assess the presence of heterogeneity, ranging from 0% to 100%. An I^2 value greater than 70% suggested obvious heterogeneity and the need for a random-effect model. An I^2 value between 40% and 70% represented moderate heterogeneity. A fixed-effect model was used for I^2 values of less than 40%, which showed that heterogeneity could be disregarded. A funnel plot or Egger's test was used to evaluate publication bias. A P -value of less than 0.05 was considered to indicate statistical significance. Subgroup analysis was performed after grouping according to the duration of treatment. Fractures were defined as fractures of any part of the body except the vertebral body. The sensitivity analysis method refers to combining the remaining studies after deleting each study in each group to




Figure 2. Study selection process flowchart.

observe whether the results are consistent with the previous ones.

Results

Search results

An initial independent search of the electronic database identified 12,573 potentially relevant articles. After careful examination of the titles, 11,989 articles were excluded based on the inclusion criteria. Of the remaining 584 articles, 512 were excluded after carefully examining the abstract, mainly due to young age, vitamin D₂ use, lack of a control group, etc. Out of the 72 articles read, 58 were excluded because of the lack of complete data results, noncompliance with the inclusion criteria listed above, and other reasons. Therefore, a total of 14 RCTs conducted between 2003 and 2022 were included in the final analysis, which contextualizes the varying follow-up durations from six months up to five years. Figure 2 provides a clear overview of the study selection process.

The main characteristics of the included studies are shown in Table 1. Eleven studies reported the concentration of 25(OH)D that was used for the main results, but a total of ten studies were included because the baseline concentrations of 25(OH)D were not provided in Trivedi's experiment [20]. The analysis of death, falls, and fractures included six, nine, and six RCTs, respectively. The follow-up period ranged from six months to five years. The average age of participants ranged from 60 to 82 years, with the concentration of baseline 25(OH)D in most RCTs being less than 75 nmol/L.

Intermittent overload doses of oral vitamin D₃ on serum 25(OH)D concentrations

Based on the observation time, subgroups analysis included three subgroups: 6 months to 1 year, 1–4 years, and longer than four years, of which the standardized mean differences (95% CI) were 1.33 (1.15, 1.52), 2.06 (1.78, 2.33), and 1.37 (1.34, 1.40), respectively (Figure 3). The heterogeneity results of the group with less than one year were moderate ($I^2 = 43\%$), and subgroup analysis was performed. To further delineate the impact of administration frequency, we analyzed a subset of studies characterized by a one-month interval between vitamin D₃ doses, termed one-month intermittent administration. This specific subgroup showed no significant heterogeneity ($I^2 = 0$). There was significant heterogeneity in the second group ($I^2 = 82\%$) and moderate heterogeneity in the one-month intermittent subgroup ($I^2 = 61\%$), possibly due to the large difference in measurement times. The funnel plot suggested that the points on both sides were asymmetric; therefore, it was analyzed for publication bias by the quantitative method of Egger's test (Egger = 0.7847), which indicated that there was no publication bias. In the sensitivity analysis, we combined the remaining study results after eliminating any study from the 1–2 years and the 1–4 years group, and it showed no change from the previous analysis, thus indicating that the results were stable.

Intermittent overload doses of oral vitamin D₃ on mortality

The RR (95% CI) for mortality for patients treated with high-dose, intermittent vitamin D compared with the control was 0.95 (0.87–1.04) (Figure 4), which was not statistically

Table 1. Characteristics of the included trials and participants

Study	Treatment	Sex, female (%)	No. of participants	Mean age (years)	Post 25(OH)D (mean)	Observation (mean time point)	Outcome
Malihi 2019 [20]	100,000 IU monthly	59.1	170	65.9	61.9 nmol/L	3.3 years	VD ^a
	Placebo	57.1	163	65.9	61.6 nmol/L		
Khaw 2017 [21]	Initial 200,000, then 100,000 IU monthly	41	2539	65.9	64 nmol/L	3.4 years	Fall Fracture
	Placebo	43	2517	65.9	63 nmol/L		
Aspray 2019 [22]	48,000 IU monthly	50.8	113	75.4	40 nmol/L	1 year	Fall VD
	12,000 IU monthly	45.2	112	74.6	40 nmol/L		
Waterhouse 2021 [23]	60,000 IU monthly	46	1045	69.3	^b	4.3 years	Fall VD
	Placebo	45.7	1048	69.3	^b		
Trivedi 2003 [24]	100,000 IU every 4 months	47.6	1345	76.1	Not mentioned	4 years	Mortality Fall Fracture
	Placebo	50	1341	75.4	Not mentioned		
Glendening 2012 [12]	150,000 IU every 3 months	100	353	76.9	65.8 nmol/L	9 months	VD Fall Fracture
	Placebo	100	333	76.5	65.8 nmol/L		
Rachel 2022 [25]	100,000 IU monthly	45.9	10662	69.3	^c	5.7 years	VD Mortality
	Placebo	45.9	10653	69.3	^c		
Sanders 2010 [26]	50,000 IU annually	100	1131	76	53 nmol/L	15 months	VD Mortality Fall Fracture
	Placebo	100	1125	76.1	45 nmol/L		
John 2017 [27]	Initial 200,000, then 100,000 IU monthly	40	71	64.5	62.1 nmol/L	1.1 years	VD
	Placebo	30	79	65.5	63.1 nmol/L		
Scragg 2017 [28]	Initial 200,000, then 100,000 IU monthly	40.9	2558	65.9	26.5 ng/mL	3.3 years	VD Mortality
	Placebo	42.9	2550	65.6	26.5 ng/mL		
Scragg 2019 [29]	initial 200,000, then 100,000 IU monthly	^d	2558	65.9	64 nmol/L	3.3 years	Fall Fracture
	Placebo	^d	2550	65.9	64 nmol/L		
Ginde 2017 [30]	100,000 IU monthly	60	55	80	23 ng/mL	1 year	Mortality Fall Fracture
	Placebo	55.8	52	82	23 ng/mL		
Rake 2020 [31]	100,000 IU monthly	53	392	^e	52.4 nmol/L	2 years	VD Mortality
	Placebo	53	395	^e	48.5 nmol/L		
Schwetz 2017 [32]	Initial 540,000, then 90,000 IU monthly	54	153	62.2	13.9 ng/mL	6 months	VD Fall Fracture
	Placebo	51	136	60	13.7 ng/mL		

^a25(OH)D concentration in serum; ^bIt was estimated that more than 76% of participants had vitamin D concentrations greater than 20 ng/mL (it was not specified precisely); ^cIt was estimated that more than 75% of participants had vitamin D concentrations greater than 20 ng/mL (it was not specified precisely); ^dThe values of the experimental group and the control group were not defined, and the average value of the two groups was 41.9%; ^eNo mean age was given, and the age range was 65 to 84; ^fFor the 25(OH)D concentration in serum, 2.5 nmol/L is equivalent to 1 ng/mL.

significant ($P = 0.25$). A total of 892 of 16,146 participants (5.5%) randomized to the vitamin D group and 937 of 16,115 participants (5.8%) randomized to the placebo or no-intervention group

died. The results remained robust after sensitivity analysis. We concluded that there was no publication bias by using Egger's test ($P = 0.7891$).




Figure 3. Subgroup analysis by observation duration for vitamin D₃ intervention. Forest plot showing standardized mean differences (SMD) for 6 months to 1 year (SMD: 1.33, 95% CI: 1.15-1.52), 1-4 years (SMD: 2.06, 95% CI: 1.78-2.33), and over 4 years (SMD: 1.37, 95% CI: 1.34-1.40). Heterogeneity was moderate to high across subgroups. Egger's test (0.7847) suggests no publication bias. CI: Confidence interval.




Figure 4. Mortality risk in patients with intermittent high-dose vitamin D₃. This forest plot illustrates the risk ratio (RR) of mortality for patients treated with intermittent high-dose vitamin D₃ versus controls. The pooled RR is 0.95 (95% CI: 0.87-1.04, $P = 0.25$), showing no significant effect on mortality. Data from 16,146 patients in the vitamin D group and 16,115 in the control group are included. CI: Confidence interval.

Intermittent overload doses of oral vitamin D₃ on falls

The RR (95% CI) for falls for patients treated with an overload dose and intermittent vitamin D compared with controls was 1.02 (0.98-1.05), without a significant difference ($P = 0.34$) (Figure 5A). The Labe diagram shows that the points were incompletely linearly distributed, and some points deviated far from the effect line, which suggested that heterogeneity was moderate for this outcome ($I^2 = 36\%$) (Figure 5B). After conducting a subgroup analysis based on the interval between drug administration, the heterogeneity disappeared ($I^2 = 0\%$) when the study by Sanders [21], which was administered intermittently for one year drug was excluded. The results remained robust after a sensitivity analysis using a funnel plot (Figure 5C) to analyze the publication bias, which suggested that the visible

points were symmetrically distributed, presenting an inverted and incomplete symmetrical funnel shape. Based on Egger's test, it was considered that there was no publication bias ($P = 0.6508$).

Intermittent overload doses of oral vitamin D₃ on fracture

The RR (95% CI) for the hip frame in patients treated with overload dose and intermittent vitamin D₃ compared with controls was 0.99 (0.84-1.18) (Figure 6), which was not statistically significant. The results suggested that intermittent overload doses of oral vitamin D₃ increased the incidence of fracture, but the sensitivity analysis showed that the CI of Sanders' experiment had changed significantly, while the statistical results and CI of other studies did not, so we deemed the results as




Figure 5. Meta-analysis forest plot assessing the impact of intermittent overload doses of oral vitamin D₃ on falls. (A) The plot compares the risk ratios (RR) for falls between experimental groups treated with vitamin D₃ and control groups across several studies. (B) The Labe plot illustrates the relationship between treatment effects and control event rates. (C) The funnel plot evaluates publication bias with symmetry around the effect size, suggesting no evidence of bias. CI: Confidence interval.




Figure 6. Forest plots of the meta-analysis of fracture. CI: Confidence interval.

unstable. However, a subsequent sensitivity analysis confirmed the stability of the results. Heterogeneity was observed for this outcome ($I^2 = 24\%$); but could be disregarded. Egger's test was used to analyze the publication bias due to limited studies, and it showed no publication bias ($P = 0.9127$).

Discussion

This meta-analysis has systematically evaluated the effects of intermittent administration of vitamin D on mortality, falls, and fractures, revealing nuanced differences between vitamin D₂ and D₃ supplements. We found that vitamin D₃, when administered intermittently, may help reduce the incidence of fractures and falls without significantly impacting mortality rates. These outcomes underscore the potential benefits of vitamin D₃ in enhancing bone health and preventing injury in at-risk

populations. A previous review also suggested that vitamin D₂ may not increase mortality [22]. However, Smith's study [23] indicated that individuals receiving vitamin D₂ had a significantly higher rate of fractures compared to the control group, raising concerns about the relative safety of this form of vitamin D. Our findings contribute to the ongoing debate by suggesting that vitamin D₂ may not offer the same level of safety as vitamin D₃, warranting further investigation into the differential impacts of these two compounds.

Decreased dose frequency has been identified as a factor associated with better responses to pharmacological therapy [24]. In addition, plasma 25(OH)D has a half-life estimated in terms of weeks rather than hours [25], so daily doses may not be required to maintain a steady vitamin D status. Most of the 25(OH)D concentrations reported by the RCTs included in this experiment were measured several

days after administration; raising doubts about whether the concentration of 25(OH)D in serum could be significantly higher in the intervention group than that in the control group after 2–3 weeks. Armas et al. [26] chose a single bolus of 50,000 IU that showed a significantly greater AUC for cholecalciferol than for ergocalciferol, with serum 25(OH)D₂ concentrations that fell rapidly back to baseline after only 14 days, whereas serum 25(OH)D₃ concentrations peaked at the same time point and had not returned to baseline for the entire 28-day intervention. Sanders [21] used an annual intermittent drug administration period and performed tests at one and three months after the initial drug administration. Although the concentration in the third month was lower than the peak concentration in the first month, it was still significantly higher than that in the control group, with a concentration greater than 75 nmol/L, which was consistent with the conclusion demonstrated by Heaney et al. [27]: large doses of the vitamin are stored in fat and then slowly converted into serum 25(OH)D. However, daily administration had more advantages in the stability of the 25(OH)D concentration in serum.

Daily, weekly, and monthly vitamin D₃ levels were compared in three trials. In one 4-month study of equivalent oral doses of vitamin D₃ (600 IU/day, 4200 IU/week, and 18000 IU/month), the daily dose was the most effective and was the only dose that increased 25(OH)D concentrations [28]. However, in another experiment with a larger sample size, the comparison of three administration methods of 1500 IU daily, 10500 IU weekly, and 45000 IU every 28 days showed the same effectiveness results across all three regimens [29]. Essentially, the mode of administration for a higher dose may have different effects compared to a lower dose. In this meta-analysis, the dose for all RCTs in the experimental group was equivalent to more than 800 IU per day, potentially achieving higher serum 25(OH)D concentrations than those expected from less frequent, higher-dose regimens. In addition, Ilahi et al. [30] suggested that the dosing interval of intermittent dosing regimens should not be greater than 70 days to ensure that 25(OH)D levels do not decline below a target concentration of 70 nmol/L. Considering the adverse results of the annual administration analyzed previously, a monthly dosing interval may be more suitable.

According to the study of the group that was observed for more than 1 year, the concentration of 25(OH)D in serum was maintained between 44 and 56 ng/mL, which was much higher than the target concentration of 30 ng/mL [31]. Subsequently, issues, such as elevated serum and urine calcium, kidney stones, and other adverse events arose. However, the majority of disease-specific recommendations state consistently that the minimum serum 25(OH)D concentration should be 30 ng/mL, with an upper limit typically ranging between 50 and 60 ng/mL. Achieving and maintaining such values require regular vitamin D supplementation with doses of 3000–5000 IU/day [32]. It is generally assumed that large doses of vitamin D₃ excreted through the kidneys can significantly increase the burden on the kidneys. Vieth et al. [33] conducted a 6-month safety and efficacy study and concluded that consumption of more than 4000 IU/day caused no harm and effectively raised 25(OH)D levels to “high-normal” concentrations (140 nmol/L)

in practically all adults. In 2011, the Institute of Medicine’s report on dietary intake of vitamin D recommended an upper limit of 4000 IU/day and stated that doses up to 10,000 IU/day were safe. The studies included in this meta-analysis did not surpass the equivalent of 10,000 IU/day [34]. Malihi’s meta-analysis [35] suggested that intermittent administration of large doses (equivalent to more than 2800 IU/day) might increase the incidence rate of high serum calcium but not the risk of high urinary calcium or kidney stones. However, the inclusion criteria for that analysis did not limit age or the method of administration of vitamin D₂ or vitamin D₃; therefore, whether this conclusion is applicable to our meta-analysis, more RCTs that meet the aforementioned conditions are needed.

Vitamin D₃ supplementation in appropriate doses is known to have a positive effect on fractures related to muscle function. However, very high doses of vitamin D can have a negative effect on muscle function due to a sudden increase in vitamin D receptor occupancy. Vitamin receptors are also present in the central nervous system [36], making it possible for falls to be affected as well. However, the exact amount of vitamin D₃ administered that causes negative neuromuscular effects is unknown. Therefore, this may explain why there was no increase in the incidence of fractures and falls in the lower interdose group between January and April (receiving less than 200,000 IU as a single dose). In contrast, Sanders’ [21] study used a dose of 500,000 IU per administration and showed a significant increase in the incidence of falls. However, it should be noted that the results of the Schwetz [37] trial with a single dose of 540,000 IU showed no significant increase in the incidence of falls and fractures. Therefore, the negative effects of larger doses need to be verified in more RCTs.

In summary, we believed that this study had no obvious publication bias due to strict criteria for selecting RCTs and reducing heterogeneity after discussing the sources of heterogeneity in the analysis. Through sensitivity analysis, we came to a clear and convincing conclusion that oral vitamin D₃ with more than 48,000 IU per month resulted in better compliance and was a more effective treatment regimen. However, this article does have some limitations. First, the intervention measures of the experimental group involved calcium, of which the preventive effect on fractures or falls was not analyzed in this article. Second, most included studies were conducted in regions far from the equator, namely, southeast Australia, New Zealand, the United Kingdom, and the United States. It is well known that the production of vitamin D is closely related to sunlight exposure, so this treatment may not be applicable to individuals living near the equator. The final conclusion cannot establish a secure upper limit for a single dose owing to insufficient evidence. However, a single dose of 200,000 IU was considered safe for administration.

Conclusions

This analysis showed that receiving a high-dose dose (equivalent to more than 800 IU per day) of oral vitamin D₃ every month for 1 year led to a significant increase in the

concentration of 25(OH)D. Test results at any time after six months were above 75 nmol/L and this did not increase the incidence of fractures, falls, and deaths. Therefore, this treatment method can be promoted in middle-aged and elderly patients in high-latitude countries. Although one-year intermittent administration significantly increased the concentration of 25(OH)D in serum, whether the method would increase the incidence of falls requires further research and, thus, is not recommended due to the lack of safety demonstration with more relevant RCTs.

Conflicts of interest: Authors declare no conflicts of interest.

Funding: Authors received no specific funding for this work.

Data availability: Please contact the corresponding author for data requests.

Submitted: 07 March 2024

Accepted: 02 April 2024

Published online: 13 April 2024

References

- [1] Wang Y, Liu Y, Lian Y, Li N, Liu H, Li G. Efficacy of high-dose supplementation with oral vitamin D3 on depressive symptoms in dialysis patients with vitamin D3 insufficiency: a prospective, randomized, double-blind study. *J Clin Psychopharmacol* 2016;36:229–35. <https://doi.org/10.1097/JCP.0000000000000486>.
- [2] Narula N, Cooray M, Anglin R, Muqtadir Z, Narula A, Marshall JK. Impact of high-dose vitamin D3 supplementation in patients with Crohn's disease in remission: a pilot randomized double-blind controlled study. *Dig Dis Sci* 2017;62:448–55. <https://doi.org/10.1007/s10620-016-4396-7>.
- [3] Hansen KE, Bartels CM, Gangnon RE, Jones AN, Gogineni J. An evaluation of high-dose vitamin D for rheumatoid arthritis. *J Clin Rheumatol* 2014;20:112–4. <https://doi.org/10.1097/RHU.0000000000000072>.
- [4] Schall JJ, Hediger ML, Zemel BS, Rutstein RM, Stallings VA. Comprehensive safety monitoring of 12-month daily 7000-IU vitamin D3 supplementation in human immunodeficiency virus-infected children and young adults. *JPEN J Parenter Enteral Nutr* 2016;40:1057–63. <https://doi.org/10.1177/0148607115593790>.
- [5] Scragg R, Sowers M, Bell C. Serum 25-hydroxyvitamin D, ethnicity, and blood pressure in the third national health and nutrition examination survey. *Am J Hypertens* 2007;20:713–9. <https://doi.org/10.1016/j.amjhyper.2007.01.017>.
- [6] Rizzoli R, Boonen S, Brandi ML, Bruyère O, Cooper C, Kanis JA, et al. Vitamin D supplementation in elderly or postmenopausal women: a 2013 update of the 2008 recommendations from the European society for clinical and economic aspects of osteoporosis and osteoarthritis (ESCEO). *Curr Med Res Opin* 2013;29:305–13. <https://doi.org/10.1185/03007995.2013.766162>.
- [7] Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab* 2011;96:1911–30. <https://doi.org/10.1210/jc.2011-0385>.
- [8] Hollis BW, Johnson D, Hulsey TC, Ebeling M, Wagner CL. Vitamin D supplementation during pregnancy: double-blind, randomized clinical trial of safety and effectiveness. *J Bone Miner Res* 2011;26:2341–57. <https://doi.org/10.1002/jbmr.463>.
- [9] Wijnen H, Saleminck D, Roovers L, Taekema D, De Boer H. Vitamin D supplementation in nursing home patients: randomized controlled trial of standard daily dose versus individualized loading dose regimen. *Drugs Aging* 2015;32:371–8. <https://doi.org/10.1007/s40266-015-0259-8>.
- [10] Dawson-Hughes B, Harris SS. High-dose vitamin D supplementation: too much of a good thing? *JAMA* 2010;303:1861–2. <https://doi.org/10.1001/jama.2010.598>.
- [11] Rossini M, Adami S, Viapiana O, Fracassi E, Idolazzi L, Povino MR, et al. Dose-dependent short-term effects of single high doses of oral vitamin D3 on bone turnover markers. *Calcif Tissue Int* 2012;91:365–9. <https://doi.org/10.1007/s00223-012-9637-y>.
- [12] Glendenning P, Zhu K, Inderjeeth C, Howat P, Lewis JR, Prince RL. Effects of three-monthly oral 150,000 IU cholecalciferol supplementation on falls, mobility, and muscle strength in older postmenopausal women: a randomized controlled trial. *J Bone Miner Res* 2012;27:170–6. <https://doi.org/10.1002/jbmr.524>.
- [13] Zheng YT, Cui QQ, Hong YM, Yao WG. A meta-analysis of high dose, intermittent vitamin D supplementation among older adults. *PLoS One* 2015;10:e0115850. <https://doi.org/10.1371/journal.pone.0115850>.
- [14] Thanapluetiwong S, Chewcharat A, Takkavatakarn K, Praditpornsilpa K, Eiam-Ong S, Susantitaphong P. Vitamin D supplement on prevention of fall and fracture: a meta-analysis of randomized controlled trials. *Medicine (Baltimore)* 2020;99:e21506. <https://doi.org/10.1097/MD.00000000000021506>.
- [15] Zhao JG, Zeng XT, Wang J, Liu L. Association between calcium or vitamin D supplementation and fracture incidence in community-dwelling older adults: a systematic review and meta-analysis. *JAMA* 2017;318:2466–82. <https://doi.org/10.1001/jama.2017.19344>.
- [16] Bischoff-Ferrari HA, Willett WC, Wong JB, Giovannucci E, Dietrich T, Dawson-Hughes B. Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *JAMA* 2005;293:2257–64. <https://doi.org/10.1001/jama.293.18.2257>.
- [17] Izaks GJ. Fracture prevention with vitamin D supplementation: considering the inconsistent results. *BMC Musculoskelet Disord* 2007;8:26. <https://doi.org/10.1186/1471-2474-8-26>.
- [18] Tripkovic L, Lambert H, Hart K, Smith CP, Bucca G, Penson S, et al. Comparison of vitamin D2 and vitamin D3 supplementation in raising serum 25-hydroxyvitamin D status: a systematic review and meta-analysis. *Am J Clin Nutr* 2012;95:1357–64. <https://doi.org/10.3945/ajcn.111.031070>.
- [19] Rothen JP, Rutishauser J, Walter PN, Hersberger KE, Arnet I. Oral intermittent vitamin D substitution: influence of pharmaceutical form and dosage frequency on medication adherence: a randomized clinical trial. *BMC Pharmacol Toxicol* 2020;21:51. <https://doi.org/10.1186/s40360-020-00430-5>.
- [20] Trivedi DP, Doll R, Khaw KT. Effect of four monthly oral vitamin D3 (cholecalciferol) supplementation on fractures and mortality in men and women living in the community: randomised double blind controlled trial. *BMJ* 2003;326:469. <https://doi.org/10.1136/bmj.326.7387.469>.
- [21] Sanders KM, Stuart AL, Williamson EJ, Simpson JA, Kotowicz MA, Young D, et al. Annual high-dose oral vitamin D and falls and fractures in older women: a randomized controlled trial. *JAMA* 2010;303:1815–22. <https://doi.org/10.1001/jama.2010.594>.
- [22] Bjelakovic G, Gluud LL, Nikolova D, Whitfield K, Wetterslev J, Simonetti RG, et al. Vitamin D supplementation for prevention of mortality in adults. *Cochrane Database Syst Rev* 2014;1:CD007470. <https://doi.org/10.1002/14651858.CD007470.pub3>.
- [23] Smith H, Anderson F, Raphael H, Maslin P, Crozier S, Cooper C. Effect of annual intramuscular vitamin D on fracture risk in elderly men and women—A population-based, randomized, double-blind, placebo-controlled trial. *Rheumatology (Oxford)* 2007;46:1852–7. <https://doi.org/10.1093/rheumatology/kem240>.
- [24] Jones KS, Assar S, Vanderschueren D, Bouillon R, Prentice A, Schoenmakers I. Predictors of 25(OH)D half-life and plasma 25(OH)D concentration in The Gambia and the UK. *Osteoporos Int* 2015;26:1137–46. <https://doi.org/10.1007/s00198-014-2905-0>.
- [25] Jin J, Sklar GE, Oh VMS, Chuen LI. Factors affecting therapeutic compliance: a review from the patient's perspective. *Ther Clin Risk Manag* 2008;4:269–86. <https://doi.org/10.2147/TCRM.S1458>.
- [26] Armas LA, Hollis BW, Heaney RP. Vitamin D2 is much less effective than vitamin D3 in humans. *J Clin Endocrinol Metab* 2004;89:5387–91. <https://doi.org/10.1210/jc.2004-0360>.
- [27] Heaney RP, Armas LA, Shary JR, Bell NH, Binkley N, Hollis BW. 25-hydroxylation of vitamin D3: relation to circulating vitamin D3 under various input conditions. *Am J Clin Nutr* 2008;87:1738–42. <https://doi.org/10.1093/ajcn/87.6.1738>.
- [28] Chel V, Wijnhoven HA, Smit JH, Ooms M, Lips P. Efficacy of different doses and time intervals of oral vitamin D

- supplementation with or without calcium in elderly nursing home residents. *Osteoporos Int* 2008;19:663–71. <https://doi.org/10.1007/s00198-007-0465-2>.
- [29] Ish-Shalom S, Segal E, Salganik T, Raz B, Bromberg IL, Vieth R. Comparison of daily, weekly, and monthly vitamin D3 in ethanol dosing protocols for two months in elderly hip fracture patients. *J Clin Endocrinol Metab* 2008;93:3430–5. <https://doi.org/10.1210/jc.2008-0241>.
- [30] Ilahi M, Armas LA, Heaney RP. Pharmacokinetics of a single, large dose of cholecalciferol. *Am J Clin Nutr* 2008;87:688–91. <https://doi.org/10.1093/ajcn/87.3.688>.
- [31] Dawson-Hughes B, Mithal A, Bonjour JP, Boonen S, Burckhardt P, Fuleihan GE, et al. IOF position statement: vitamin D recommendations for older adults. *Osteoporos Int* 2010;21:1151–4. <https://doi.org/10.1007/s00198-010-1285-3>.
- [32] Ekwaru JP, Zwicker JD, Holick MF, Giovannucci E, Veugelers PJ. The importance of body weight for the dose response relationship of oral vitamin D supplementation and serum 25-hydroxyvitamin D in healthy volunteers. *PLoS One* 2014;9:e111265. <https://doi.org/10.1371/journal.pone.0111265>.
- [33] Vieth R, Chan PC, MacFarlane GD. Efficacy and safety of vitamin D3 intake exceeding the lowest observed adverse effect level. *Am J Clin Nutr* 2001;73:288–94. <https://doi.org/10.1093/ajcn/73.2.288>.
- [34] Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM, Clinton SK, et al. The 2011 report on dietary reference intakes for calcium and vitamin D from the institute of medicine: what clinicians need to know. *J Clin Endocrinol Metab* 2011;96:53–8. <https://doi.org/10.1210/jc.2010-2704>.
- [35] Malihi Z, Wu Z, Lawes CMM, Scragg R. Adverse events from large dose vitamin D supplementation taken for one year or longer. *J Steroid Biochem Mol Biol* 2019;188:29–37. <https://doi.org/10.1016/j.jsbmb.2018.12.002>.
- [36] Buell JS, Dawson-Hughes B. Vitamin D and neurocognitive dysfunction: preventing “D” ecliptic? *Mol Aspects Med* 2008;29:415–22. <https://doi.org/10.1016/j.mam.2008.05.001>.
- [37] Schwetz V, Schnedl C, Urbanic-Purkart T, Trummer C, Dimai HP, Fahrleitner-Pammer A, et al. Effect of vitamin D3 on bone turnover markers in critical illness: post hoc analysis from the VITDAL-ICU study. *Osteoporos Int* 2017;28:3347–54. <https://doi.org/10.1007/s00198-017-4190-1>.

Related articles published in BJBMS

1. Vitamin D status, serum lipid concentrations, and vitamin D receptor (VDR) gene polymorphisms in familial mediterranean fever
Turan Turhan et al., BJBMS, 2018