

The Immunostimulatory Properties of Vitamin D

Khalil Ismail Abid Mohammed^{1*}, Sawsan H. Authman², Sarmad M. Zeiny³, Saad Hasan Mohammed Ali⁴

¹Clinical Communicable Disease Research Unit /College of Medicine, University of Baghdad, Baghdad, Iraq

²Department of Biology, College of Science, AL-Mustansiriyah University, Baghdad, Iraq

³Department of Microbiology, College of Medicine, University of Baghdad, Baghdad, Iraq

⁴Al-Mustaqbal University College of Dentistry

DOI: <https://doi.org/10.36348/sjls.2025.v10i05.004>

| Received: 28.03.2025 | Accepted: 05.05.2025 | Published: 13.05.2025

*Corresponding author: Khalil Ismail Abid Mohammed

Clinical Communicable Disease Research Unit /College of Medicine, University of Baghdad, Baghdad, Iraq

Abstract

Vitamin D has been associated with the regulation of bone metabolism. However, increasing evidence demonstrates a strong association between vitamin D signaling and many biological processes to that immunostimulatory and immunomodulatory effects on the immune system. the presence of the vitamin D receptor in multiple immune cells, such as monocytes, dendritic cells, and activated T cells, vitamin D with a novel role in modulating immunological functions and its subsequent role in the development or prevention of autoimmune diseases such as systemic lupus Erythematosus, Type-1 Diabetes mellitus and multiple sclerosis. In the present review, the multiple and diverse effects of vitamin D on the immune system are reviewed.

Keywords: Vitamin D, Immune system, Immunomodulation, Vitamin D receptor (VDR), Vitamin D receptor (VDR).

Copyright © 2025 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

The name vitamin D covers a group of liposoluble steroid compounds of different origins with

similar chemical structures and the same biological effects. Two main forms of vitamin D are ergocalciferol (vitamin D 2) and cholecalciferol (vitamin D 3) (Figure 1).




Figure 1: Chemical structure of vitamin D2 (A), vitamin D3 (B), and vitamin D4 (C). In online version, differences are shown in blue

Vitamin D 1 is a historical term for a mixture of vitamin D2 with lumisterol. Vitamin D 2 is Synthesized by the irradiation of ergosterol in yeast, while vitamin D

3 is generated from 7-dehydrocholesterol after ultraviolet (UV)-B irradiation in the human skin this being a unique property among vitamins (Figure 2).




Figure 2: Synthesis of vitamin D

Although vitamin D 2 and D3 were considered equally active for many years, current knowledge indicates that the potency of vitamin D2 is less than one-third of that of vitamin D3 [1-3]. The potential responsible factors are different metabolic pathways and/or different affinity of the active metabolites of vitamins D2 and D3 toward vitamin D receptor (VDR). Vitamin D3 is the main form of vitamin D in humans, and indeed, it is estimated that about 80–90% of the vitamin D requirements are covered by the endogenous synthesis in the extent of the skin vitamin D synthesis is dependent on the length of skin sun exposure, the season of the year, and latitude [4]. A 20-min long whole-body exposure to the summer sun is able to produce up to 250 lg of vitamin D3 [5, 6]. A number of effects are attributed to vitamin D. The most known and studied effect is linked to calcium and phosphate homeostasis, with a crucial impact on bone metabolism. In addition to mineral homeostasis, current research has been investigating a plethora of different activities of vitamin D. VDR, which interacts with the active form of vitamin D, calcitriol or 1,25-dihydroxy-cholecalciferol

(1,25(OH)₂D₃) has been identified in almost all human cells, even in those that do not play a role in calcium metabolism, including the skin, brain, immune cells, prostate cancer cells, and pancreases [7, 8].

Vitamin D and Immunity

While it is well established that vitamin D enhances intestinal calcium absorption, an effect mediated via regulation of calcium transport proteins in the small intestine [9], exhibiting a central role in its effects on the immune system (Figure 3).

Cells of the immune system harbour the vitamin D activating enzyme 1- α -hydroxylase and express the vitamin D receptor (VDR) [10, 11]. Cells of the immune system which express the VDR and harbour 1- α -hydroxylase are macrophages, T cells, dendritic cells, monocytes, and B cells [12-14] (Figure -4).

Vitamin D is involved both in the regulation of the innate immunity as it enhances the body defense system.




Figure 3: Cells of the immune system which are targets of vitamin D, macrophages, neutrophils, T-lymphocytes, dendritic cells, B lymphocytes




Figure 4: The effects of vitamin D on the immune system

Against microbes and other pathogenic organisms, as well as in the modulation of the adaptive immune system through direct effects on T cell activation and on antigen presenting cells; in particular, dendritic cells.

Vitamin D and the Innate Immune System

Vitamin D regulates the innate immune system [15-17]. The innate immune system - strategy is a first

line of defense against infection [18, 19]. Its major functions include recruitment of immune cells, activation of the complement cascade (Figure -5). Identification and removal of foreign substances, activation of the adaptive immune response, and the utilization of physical and chemical barriers against infectious agents [19]. The vitamin D receptor (VDR) is expressed both in the keratinocytes [20, 21], and cells of the innate immune system such as macrophages and monocytes [22-25].




Figure 5: Vitamin D and innate immunity

Vitamin D enhances the production of defensin $\beta 2$ and cathelicidin in response to infection by macrophages, monocytes, and keratinocytes [26]. Humans have only one cathelicidin [27], which is produced by cells of the immune system, including neutrophils, macrophages, and cells lining epithelial surfaces that are constantly exposed to potential pathogens such as the skin, the respiratory, and the gastrointestinal tract [28-30].

Vitamin D and Adaptive Immunity

Vitamin D acts to regulate the adaptive immune system (Figure-6). The adaptive immune system includes both humeral immunity components and cell mediated immunity components, both directed against invading pathogens. Adaptive immunity leads to immunological memory after an initial response to a specific pathogen, resulting in an enhanced response to future encounters with that pathogen [31]. Vitamin D can promote development and function of Tregs in vitro. Also, effector T cells are directly and indirectly affected

leading to a shift in the Th1/Th2 balance toward Th2 and a reduction of the Th17 response [32]. Once T cells are activated, 1,25(OH) $_2$ D3 inhibits IL-2 production [33]. T cells harbour the vitamin D receptor. The behaviour of T cells is modulated by vitamin D indirectly via its effects on dendritic cells. The vitamin D receptor is expressed at low levels in freshly isolated CD8 $^+$ and CD4 $^+$ T cells [34, 35], vitamin D deficiency results in a reduced capacity to turn off T cells following activation. (36) In a previous study, peripheral blood mononuclear cells which were stimulated with T-cell specific mitogen in the presence of 1,25(OH) $_2$ D3 proliferated less and produced less inflammatory cytokines, including interferon- γ [37].

B cells can upregulate the expression of vitamin D receptor and 1 α -hydroxylase [38]. 1,25(OH) $_2$ D3 in B cells can induce apoptosis, inhibiting memory B cell formation and preventing differentiation of B cells to immunoglobulin-producing plasma cells [39].




Figure 6: Vitamin D and Adaptive immunity

Vitamin D and Autoimmune Disease

Multiple Sclerosis (MS)

MS is a chronic inflammatory disease characterized by immune-mediated damage of central nervous system [40]. The etiology of MS is not well understood but, like other autoimmune diseases, could be attributed to genetic predisposition and/or environmental factors [41, 42], the prevalence of the disease is lower in equatorial regions and becomes higher with increasing latitudes [43]. This phenomenon could be caused by the lack of sunshine in high-latitude regions that is required for the cutaneous synthesis of vitamin D and suggests that vitamin D insufficiency could be a potential risk factor for MS. Multiple studies also demonstrated that most of the MS patients were deficient in vitamin D [44-46]. In particular, low serum 25 (OH)D levels were associated with high disability and relapse rate in MS patients [44, 45]. However, serum 1,25 (OH)₂D, which is the biologically active form of vitamin D, was not directly associated with both disability and relapse rate [45]. Recent studies showed that vitamin D deficiency might increase the risk of MS [47]. As a result, detection of vitamin D deficiency and restoring vitamin D status to adequate levels the clinical trials were unable to provide definitive evidence to support the therapeutic effects of vitamin D intervention, the beneficial effects of oral vitamin D supplementation on MS progression need to be further investigated [48].

Type 1 Diabetes Mellitus (T1DM)

T1DM is an immune-mediated disease that is caused by autoimmune destruction of pancreatic β cells that produce insulin, leading to insulin deficiency [49].

Similar to other autoimmune diseases, environmental factors and/or genetic susceptibility could play roles in the onset and progression of T1DM [50]. The prevalence of vitamin D deficiency/insufficiency was higher in T1DM children compared to controls [51-53]. The strong correlation between vitamin D and T1DM, it has been suggested that vitamin D could prevent the damage, rescue the function of pancreatic β cells, and reduce the incidence of T1DM. Pitocco *et al.*, and Li *et al.*, both demonstrated that vitamin D had a protective effect on preserving β -cell function [54, 55].

Systemic Lupus Erythematosus (SLE)

SLE is a systemic autoimmune disease that can cause chronic inflammation and damage in multiple tissues and organs [56]. Environmental factors and genetic susceptibility are both responsible for the pathogenesis of SLE [57, 58]. One of such factors is vitamin D deficiency. SLE patients tend to have inadequate vitamin D since most of them are photosensitive to UV radiation and unable to expose themselves to sunlight [59]. The correlation between vitamin D deficiency/insufficiency and SLE has been documented in multiple studies. Some studies suggested that vitamin D concentrations were inversely related to SLE activity [60-62].

Rheumatoid Arthritis

Vitamin D does not play an important role in the pathogenesis of rheumatoid arthritis. This fact was supported by a recent studies, which showed that there are insufficient data to demonstrate the effect of vitamin D on rheumatoid arthritis activity (DAS-28 or pain) or

flares [63]. On the contrary, vitamin D supplementation can be employed to prevent musculoskeletal complications in patients suffering from rheumatoid arthritis with a deficiency of this vitamin.

CONCLUSION

Vitamin D is a likely immunostimulatory or immunomodulatory agent. It has immune stimulating properties, as it enhances the function of the innate and Adaptive immune system. Vitamin D deficiency or insufficiency is associated with compromised immunity, leading to increased infectious diseases such as tuberculosis, and increased susceptibility to autoimmune diseases such as type 1 diabetes, Rheumatoid arthritis, systemic Lupus Erythematosus and others.

Conflicts of Interest: The authors declare no conflicts of interest.

REFERENCES

- Armas LA, Hollis BW, Heaney RP. Vitamin D2 is muchless effective than vitamin D3 in humans. *J Clin Endocrinol Metab*. 2004;89(11):5387–5391.
- Tjellesen L, Christiansen C, Rodbro P, et al. Different metabolism of vitamin D2 and vitamin D3 in epileptic patients on carbamazepine. *Acta Neurol Scand*.1985;71(5):385–389.
- Tjellesen L, Gotfredsen A, Christiansen C. Different actions of vitamin D2 and D3 on bone metabolism in patients treated with phenobarbitone/phenytoin. *Calcif Tissue Int*. 1985;37(3):218–222.
- Borel P, Caillaud D, Cano N. Vitamin D bioavailability: state of the art. *Crit Rev Food Sci Nutr*. 2015;55(9):1193–1205
- Krause R, Böhning M, Hopfenmüller W, et al. Ultraviolet B and blood pressure. *Lancet*. 1998; 352(9129):709–710.
- Stamp T, Haddad J, Twigg C. Comparison of oral 25-hydroxycholecalciferol, vitamin D, and ultraviolet light as determinants of circulating 25-hydroxyvitamin D. *Lancet*. 1977;1(8026):1341–1343.
- Wang Y, Zhu J, DeLuca HF. Where is the vitamin D receptor? *Arch Biochem Biophys*. 2012;523(1): 123–133.
- Christakos S, DeLuca HF. Minireview: vitamin D: is there a role in extra skeletal health ?*Endocrinology*.2011;152(8):2930–2936
- Wacker M, Holick MF. Vitamin D - effects on skeletal and extraskeletal health and the need for supplementation. *Nutrients*2013;5(1):111-48. 10-
- Morán-Auth Y, Penna-Martinez M, Shoghi F, Ramos-Lopez E,Badenhoop K. Vitamin D status and gene transcription in immune cells. *J Steroid Biochem Mol Biol* 2013;136:83-5.
- Szymczak I, Pawliczak R. The Active Metabolite of Vitamin D3 as a Potential Immunomodulator. *Scand J Immunol* 2016;83(2):83-91.
- Veldman CM, Cantorna MT, DeLuca HF. Expression of 1,25-dihydroxyvitamin D(3) receptor in the immune system. *Arch BiochemBiophys* 2000;374(2):334-338.
- Froicu M, Cantorna MT. Vitamin D and the vitamin D receptor arecritical for control of the innate immune response to colonic injury.*BMC Immunol* 2007;8:5.
- Takahashi K, Nakayama Y, Horiuchi H, Ohta T, Komoriya K, Ohmori H, et al. Human neutrophils express messenger RNA of vitamin D receptor and respond to 1alpha,25-dihydroxyvitaminD3. *Immunopharmacol Immunotoxicol* 2002;24(3):335-47.
- Christakos S, Li S, De La Cruz J, Bikle DD. New developments in our understanding of vitamin metabolism, action and treatment .*Metabolism* 2019;98:112-20
- Wei R, Christakos S. Mechanisms Underlying the Regulation of Innate and Adaptive Immunity by Vitamin D. *Nutrients* 2015;7(10):8251-60
- Jaime J, Vargas-Bermúdez DS, Yitbarek A, Reyes J,RodríguezLecompteJC. Differentialimmunomodulatory effect of vitamin D (1,25 (OH)(2) D(3)) on the innate immune response in different types of cells infected in vitro with infectious bursal disease virus.*Poult Sci* 2020;99(9):4265-77
- Hato T, Dagher PC. How the Innate Immune SystemSenses Trouble and Causes Trouble. *Clin J Am Soc Nephrol*2015;10(8):1459-69.
- McComb S, Thiriot A, Akache B, Krishnan L, Stark F. Introductionto the Immune System. *Methods Mol Biol* 2019;2024:1-24.
- Ge X, Wang L, Li M, Xu N, Yu F, Yang F, et al. Vitamin D/VDRsignaling inhibits LPS-induced IFN γ and IL-1 β in Oral epitheliaby regulating hypoxia-inducible factor-1 α signaling pathway. *CellCommun Signal* 2019;17(1):18.
- Bikle DD. The Vitamin D Receptor as Tumor Suppressor in Skin.*Adv Exp Med Biol*2020;1268:285-306. 56.
- 22-Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, et al.Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 2006;311(5768):1770-3.
- Martens PJ, Gysemans C, Verstuyf A, Mathieu AC. Vitamin D'sEffect on Immune Function. *Nutrients* 2020;12(5).
- Baeke F, Takiishi T, Korf H, Gysemans C, Mathieu C. VitaminD: modulator of the immune system. *Curr Opin Pharmacol*2010;10(4):482-96.
- Kloc M, Ghobrial RM, Lipińska-Opalka A, Wawrzyniak A,Zdanowski R, Kalicki B, et al. Effects of vitamin D on macrophages and myeloid-derived suppressor cells (MDSCs) hyper inflammatory response in the lungs of COVID-19 patients. *Cell Immunol* 2021;360:104259.
- Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, et al.Toll-like receptor triggering of a vitamin

- D-mediated human antimicrobial response. *Science* 2006;311(5768):1770-3.
27. Xhindoli D, Pacor S, Benincasa M, Scocchi M, Gennaro R, Tossi A. The human cathelicidin LL-37-A pore-forming antibacterial peptide and host-cell modulator. *Biochim Biophys Acta* 2016;1858(3):546-66
28. Weber G, Heilborn JD, Chamorro Jimenez CI, Hammarsjo A, Torma H, Stahle M. Vitamin D induces the antimicrobial protein hCAP18 in human skin. *J Invest Dermatol* 2005 May;124(5):1080-2.
29. Bals R, Wang X, Zasloff M, Wilson JM. The peptide antibiotic LL-37/hCAP-18 is expressed in epithelia of the human lung where it has broad antimicrobial activity at the airway surface. *Proc Natl Acad Sci U S A* 1998;95(16):9541-6.
30. Gallo RL, Kim KJ, Bernfield M, Kozak CA, Zanetti M, Merluzzi L, et al. Identification of CRAMP, a cathelin-related antimicrobial peptide expressed in the embryonic and adult mouse. *J Biol Chem* 1997;272(20):13088-93.
31. Shampo MA, Kyle RA. Adolf Windaus--Nobel Prize for research on sterols. *Mayo Clin Proc* 2001;76(2):119
32. Peelen E, Knippenberg S, Muris AH, Thewissen M, Smolders J, Tervaert JW, et al. Effects of vitamin D on the peripheral adaptive immune system: a review. *Autoimmun Rev* 2011;10(12):733-43.
33. Chen J, Bruce D, Cantorna MT. Vitamin D receptor expression controls proliferation of naïve CD8⁺ T cells and development of CD8 mediated gastrointestinal inflammation. *BMC Immunol* 2014;15:6
34. Provvedini DM, Tsoukas CD, Deftos LJ, Manolagas SC. 1,25-dihydroxyvitamin D₃ receptors in human leukocytes. *Science* 1983;221(4616):1181-3.
35. Manolagas SC, Provvedini DM, Tsoukas CD. Interactions of 1,25-dihydroxyvitamin D₃ and the immune system. *Mol Cell Endocrinol* 1985;43(2-3):113-22.
36. Ooi JH, Chen J, Cantorna MT. Vitamin D regulation of immune function in the gut: why do T cells have vitamin D receptors? *Mol Aspects Med* 2012;33(1):77-82.
37. Rigby WF, Denome S, Fanger MW. Regulation of lymphokine production and human T lymphocyte activation by 1,25-dihydroxyvitamin D₃. Specific inhibition at the level of messenger RNA. *J Clin Invest* 1987;79(6):1659-64.
38. Chen S, Sims GP, Chen XX, Gu YY, Lipsky PE. Modulatory effects of 1,25-dihydroxyvitamin D₃ on human B cell differentiation. *J Immunol* 2007;179(3):1634-47.
39. Lemire JM, Adams JS, Sakai R, Jordan SC. 1 α ,25-dihydroxyvitamin D₃ suppresses proliferation and immunoglobulin production by normal human peripheral blood mononuclear cells. *J Clin Invest* 1984;74(2):657-61
40. Broux B, Pannemans K, Zhang X, et al. CX(3)CR1 drives cytotoxic CD4(+)CD28(-) T cells into the brain of multiple sclerosis patients. *J Autoimmun.* 2012; 38:10–19.
41. Luckey D, Bastakoty D, Mangalam AK. Role of HLA class II genes in susceptibility and resistance to multiple sclerosis: studies using HLA transgenic mice. *J Autoimmun.* 2011; 37:122–128.
42. Moroni L, Bianchi I, Lleo A. Geoepidemiology, gender and autoimmune disease. *Autoimmun Rev.* 2012; 11:A386–A392.
43. Sellner J, Kraus J, Awad A, Milo R, Hemmer B, Stuve O. The increasing incidence and prevalence of female multiple sclerosis—a critical analysis of potential environmental factors. *Autoimmun Rev.* 2011; 10:495–502.
44. Van der Mei IA, Ponsonby AL, Dwyer T, et al. Vitamin D levels in people with multiple sclerosis and community controls in Tasmania, Australia. *J Neurol.* 2007; 254:581–590.
45. Smolders J, Menheere P, Kessels A, Damoiseaux J, Hupperts R. Association of vitamin D metabolite levels with relapse rate and disability in multiple sclerosis. *Mult Scler.* 2008; 14:1220–1224.
46. Hiremath GS, Cettomai D, Baynes M, et al. Vitamin D status and effect of low-dose cholecalciferol and high-dose ergocalciferol supplementation in multiple sclerosis. *Mult Scler.* 2009; 15:735–740
47. Cantorna MT. Vitamin D and its role in immunology: multiple sclerosis, and inflammatory bowel disease. *Prog Biophys Mol Biol.* 2006; 92:60–64.
48. Antico A, Tampoia M, Tozzoli R, Bizzaro N. Can supplementation with vitamin D reduce the risk or modify the course of autoimmune diseases? A systematic review of the literature. *Autoimmun Rev.* 2012; 12:127–136.
49. Roy E, Leduc M, Guegan S, et al. Specific maternal micro-chimeric T cells targeting fetal antigens in beta cells predispose to autoimmune diabetes in the child. *J Autoimmun.* 2011; 36:253–262
50. Bogdanos DP, Smyk DS, Rigopoulou EI, et al. Twin studies in autoimmune disease: genetics, gender and environment. *J Autoimmun.* 2012; 38:J156–J169.
51. Borkar VV, Devidayal, Verma S, Bhalla AK. Low levels of vitamin D in North Indian children with newly diagnosed type 1 diabetes. *Pediatr Diabetes.* 2010; 11:345–350.
52. Singh DK, Winocour P, Summerhayes B, Viljoen A, Sivakumar G, Farrington K. Are low erythropoietin and 1,25-dihydroxyvitamin D levels indicative of tubulo-interstitial dysfunction in diabetes without persistent microalbuminuria? *Diabetes Res Clin Pract.* 2009; 85:258–264.
53. Bener A, Alsaied A, Al-Ali M, et al. High prevalence of vitamin D deficiency in type 1 diabetes mellitus and healthy children. *Acta Diabetol.* 2009; 46:183–189.
54. Li X, Liao L, Yan X, et al. Protective effects of 1 α -hydroxyvitamin D₃ on residual beta-cell function in patients with adult-onset latent

- autoimmune diabetes (LADA). *Diabetes Metab Res Rev.* 2009; 25:411–416.
55. Pitocco D, Crino A, Di Stasio E, et al. The effects of calcitriol and nicotinamide on residual pancreatic beta-cell function in patients with recent-onset type 1 diabetes (IMDIAB XI). *Diabet Med.* 2006; 23:920–923.
56. Agmon-Levin N, Mosca M, Petri M, Shoenfeld Y. Systemic lupus erythematosus one disease or many? *Autoimmun Rev.* 2012; 11:593–595.
57. Borchers AT, Naguwa SM, Shoenfeld Y, Gershwin ME. The geoepidemiology of systemic lupus erythematosus. *Autoimmun Rev.* 2010; 9:A277–A287.
58. Fu SM, Deshmukh US, Gaskin F. Pathogenesis of systemic lupus erythematosus revisited 2011: end organ resistance to damage, autoantibody initiation and diversification, and HLA-DR. *J Autoimmun.* 2011; 37:104–112.
59. Lehmann P, Homey B. Clinic and pathophysiology of photosensitivity in lupus Erythematosus. *Autoimmun Rev.* 2009; 8:456–461.
60. Amital H, Szekanecz Z, Szucs G, et al. Serum concentrations of 25-OH vitamin D in patients with systemic lupus erythematosus (SLE) are inversely related to disease activity: is it time to routinely supplement patients with SLE with vitamin D? *Ann Rheum Dis.* 2010; 69:1155–1157.
61. Wright TB, Shults J, Leonard MB, Zemel BS, Burnham JM. Hypo vitaminosis D is associated with greater body mass index and disease activity in pediatric systemic lupus erythematosus. *J Pediatr.* 2009; 155:260–265.
62. Borba VZ, Vieira JG, Kasamatsu T, Radominski SC, Sato EI, Lazaretti-Castro M. Vitamin D deficiency in patients with active systemic lupus erythematosus. *Osteoporos Int.* 2009; 20:427–433.
63. Nguyen Y, Sigaux J, Letarouilly J-G, et al. Efficacy of oral vitamin supplementation in inflammatoryrheumatic disorders: a systematic review and metaanalysis of randomized controlled trials. *Nutrients.* 2020;13(1):107.